

A review of pig and poultry diseases in the Eastern Cape Province of South Africa, 2000–2020

V Simbizi,^{1,2} R Moerane,² G Ramsay,³ C Mubamba,⁴ C Abolnik,² B Gummow^{2,5}

¹Department of Rural Development and Agrarian Reform, State Veterinary Services, South Africa

²Department of Production Animal Studies, Faculty of Veterinary Science, University of Pretoria, South Africa

³School of Animal & Veterinary Sciences and Graham Centre for Agricultural Innovation, Charles Sturt University, Australia

⁴Department of Veterinary Services, Ministry of Livestock and Fisheries, Zambia

⁵Discipline of Veterinary Sciences, College of Public Health, Medical and Veterinary Sciences, James Cook University, Australia

Corresponding author, email: vsimbizi@gmail.com

The informal poultry and pig sector in the Eastern Cape Province (ECP) of South Africa is of significant socio-economic importance as it sustains livelihoods and ensures food security; yet little is known about the distribution and prevalence of infectious and zoonotic diseases in this region. This paper reviews data published for pig and poultry diseases in the province during the last 20 years (2000–2020). The review included relevant published papers identified by a computerised literature search from Web of Science; provincial animal health reports; the national database from the Department of Agriculture, Land Reform and Rural Development (DALRRD); animal health reports submitted by DALRRD to the World Organisation for Animal Health (OIE) via the World Animal Health Information Database (WAHID) interface and laboratory records. A publication was considered eligible if it included qualitative or quantitative information on any disease affecting pigs and poultry including zoonosis. The search retrieved 174 publications, of which 26 were relevant. The review found that Newcastle disease (ND), coccidiosis and fowl pox (FP) were the most reported avian diseases in the national database, whereas avian infectious bronchitis (AIB), ND and highly pathogenic avian influenza (HPAI) were the most reported diseases in the OIE database. Classical swine fever (CSF) was the most reported pig disease in both databases. The retrieved literature on pig and poultry diseases was scarce and no longer up to date, providing decision makers with little information. The review identified important zoonotic diseases that require further studies yet failed to find information on important neglected diseases like leptospirosis.

Keywords: pig, poultry, diseases, zoonotic, Eastern Cape Province, review

Introduction

Transboundary animal diseases are highly contagious epidemic diseases that can spread extremely rapidly, irrespective of national borders. They cause mortality and morbidity in animals, thereby having serious socio-economic and sometimes public health consequences (FAO 2020). The Eastern Cape is the second largest province in South Africa after Northern Cape (Figure 1). It is divided into two metropolitan municipalities and six district municipalities. The district municipalities are in turn divided into 27 local municipalities. The human population is estimated to be 6 734 001 (STATS 2020) with the density of 39/km². The main industries include agriculture and mining (primary sector), which contribute 2% to the provincial GDP; manufacturing, electricity and construction (secondary sector) contributing 18.5% to the GDP; trade, transport, finance, personal services and government services (tertiary sector) contributing 79.5% to the GDP (ECSECC 2018). Overall the province only contributes 8% to the national GDP (STATS 2018). The Eastern Cape Province (ECP) is economically the poorest province in South Africa where subsistence agriculture predominates in the former homelands.

Livestock plays a major role in the social, cultural and economic environment in the province. The Eastern Cape is among the lowest pork and poultry producing provinces with 6% and 6.5% of total production countrywide respectively (DAFF 2018; SAPA 2017). These production statistics are mainly commercial and

do not include backyard chickens (indigenous chickens) and free roaming pigs owned by many households in the province. The informal pig and poultry sector in the ECP is estimated to have 3 841 174 birds and 536 108 pigs (STATS 2016). Apart from being a source of income for many households, pigs and poultry constitute a cheap source of protein for rural communities and ensures food security.

However, little has been published on what diseases are present in these animals within the province. Due to financial constraints, animal disease detection in the province is mainly dependent on passive surveillance in village communities (Fisher 2018, personal communication). This constitutes a major challenge since some diseases are being underreported or are not reported. Also, the province does not have animal health information systems which could help for the collection and analysis of animal health data. Such animal health information is recognised as necessary for the setting of animal health priorities (Morris 1991). Therefore, a systematic review of peer-reviewed articles, animal health reports and laboratory records compiling information on pig and poultry diseases in the province is presented with the view of identifying diseases of pigs and poultry kept within these rural communities. This will help decision makers to prioritise resources for animal disease surveillance and control in these communities once animal health information is available.




Figure 1: (a) Map of Eastern Cape Province; (b) Eastern Cape Province with its municipalities

Methods

A review was carried out on what has been published on diseases of pigs and poultry in the ECP over the last 20 years (2000–2020). The review included relevant published papers identified by a computerised literature search of all databases (WOS, BCI, CABI, CCC, DRCI, DIIDW, FSTA, KJD, MEDLINE, RSCI, SciELO and ZOOREC) from Web of Science (Appendix 7), which is the global standard for finding and connecting scholarly content across multiple disciplines around the world; monthly reports on the animal health situation submitted by the Directorate of Veterinary Services in the province to the Department of Agriculture, Land Reform and Rural Development (DALRRD); the national database from DALRRD; official animal health reports submitted by DALRRD to the World Organisation for Animal Health (OIE) and laboratory records from three provincial laboratories (Grahamstown, Middleburg and Queenstown).

Search strategy

All databases from Web of Science

All databases mentioned above were searched for published articles on pig diseases in the province from 2000 to 2020 using the following key words: Pigs OR Pig OR Swine OR Porcine (Search 1); Diseases (Search 2) and “Eastern Cape” OR (east* AND cape*) (Search 3).

Search 1, search 2 and search 3 were combined and all the published papers relevant to pig diseases in the ECP were selected.

The same search strategy was used for poultry and all databases were searched for published articles on poultry diseases in the province from 2000 to 2020 using the following key words: Chickens OR Chicken OR Poultry (Search 1); Diseases (Search 2) and “Eastern Cape” OR (east* AND cape*) (Search 3).

National database from DALRRD

The national database from DALRRD comprises all the disease reports from each province in South Africa. Each province consolidates different disease reports from the state veterinarians on a monthly basis. The Animal Diseases Act (35 of 1984) requires that all occurrences of controlled and notifiable diseases be reported to the national directorate. For other diseases and

vaccinations, the national directorate requests provinces to include them in monthly reports for OIE reporting purposes and to serve as indication of the presence and prevalence of these diseases. Some diseases that are not controlled can still have trade implications (DAFF 2016). The final report from each province is then submitted to the epidemiology section of DALRRD, which in turn, compiles and updates its national database. All disease reports from the ECP were reviewed from 1999 to 2019. The national database comprises diseases that were reported from 1993 to 2019.

WAHID interface

All official animal health reports submitted by DALRRD to the World Organisation for Animal Health (OIE) were reviewed via the World Animal Health Information Database (WAHID) interface (OIE 2020b) from 2005 to 2020.

Laboratory records at three provincial laboratories

Laboratory records were used to select pig and poultry diseases that were diagnosed at each of the three provincial laboratories in the province (Queenstown, Middleburg and Grahamstown).

Eligibility criteria

Inclusion criteria

A publication was considered eligible for this review if it included qualitative or quantitative information on any disease (bacterial, viral, parasitic and fungal) affecting pigs or poultry in the ECP. To have a wide range of reported diseases in the province, diseases affecting pigs or poultry from commercial farms were also included. Diseases affecting “poultry” other than chickens were also included. Finally, zoonotic diseases were also included in this review.

Exclusion criteria

Duplicate articles were excluded. Different references from the same study were counted as one reference irrespective of the format in which they were published (article, proceedings, workshop, etc.). The inclusion and exclusion criteria were applied to the title and abstract of all retrieved references.

Data collection process

The data collection process was undertaken in two steps. First, basic information was collected from all retrieved articles to assess which diseases have been reported in the province. For this basic analysis, the following information was systematically recorded: the publication date, the district, the species, the disease, the type of causative agent (bacteria, virus, parasite, alga, toxins, tumour, fungi, etc.), whether or not the reference focus was of a zoonotic disease, and the type of study (case report, case series, review or survey). In a second step, considering that the objective of this review was to obtain a better understanding of the current pig and poultry disease situation in the ECP, only documents published or written in the last 20 years were selected to focus on the most recent information. A more detailed analysis of the key findings from these references was then performed. The number of reported outbreaks for each disease was used to determine which disease was more frequently reported than others.

Results

Selected references and characteristics

The search strategy retrieved 174 publications, of which 26 were relevant based on the inclusion and exclusion criteria (Table I and Table II). Eighteen references were surveys (69.2%), four were case reports (15.4%), one was a conference paper

(3.8%) and three were general papers describing a particular disease nationally with little data provided for the ECP (11.5%). The majority of references provided data on diseases for pigs (84.6%), whereas references for poultry represented 15.4%. A paper on both chicken and pig disease was represented by three references (11.5%). Seventy-seven per cent of the references referred to zoonotic diseases. The following zoonotic diseases (or agents) were found in this review: Hepatitis E virus; *Enterococcus*, *Salmonella*, *E. coli*, cysticercosis, chlamydiosis, campylobacteriosis, norovirus, avian influenza, Newcastle and nocardiosis (Appendix 1).

Selected diseases from national database, OIE and laboratories records

A total of 14 diseases (10 poultry diseases and four pig diseases) were retrieved from the national database (Table III). Poultry diseases were subdivided into three categories: viral, bacterial and protozoal diseases (Figure 3). Viral diseases were most often reported (135 reported outbreaks representing 73% of all the outbreaks) followed by protozoal diseases (37 outbreaks; 20%) and bacterial diseases (13 outbreaks; 7%) (Figure 3). Among viral diseases, Newcastle disease (ND) was the most reported disease in the ECP with 103 outbreaks in the past 20 years followed by fowl pox (FP) with 18 outbreaks; avian leukosis (AL) with nine outbreaks; Gumboro and avian infectious bronchitis (AIB) with two outbreaks each and avian infectious laryngotracheitis (AIL) with one outbreak (Figure 2).

Table I: Pig diseases identified in the Eastern Cape Province between 2000 and 2020 from all databases from Web of Science

Disease	District	Year	Reference
Campylobacteriosis*	OR Tambo	2020	Ngobese et al. 2020
Campylobacteriosis*	Chris Hani and Amathole	2020	Igwaran & Okoh 2020
Classical swine fever	Eastern Cape**	2010	Akol & Lubisi 2010
<i>E. coli</i>	Amathole	2016	Iwu et al. 2016b
<i>E. coli</i>	Amathole	2017	Iwu et al. 2017
<i>Enterococcus</i>	Amathole	2015	Iweriebor et al. 2015
Hepatitis E virus	Chris Hani and Amathole	2017	Adelabu et al. 2017
Norovirus	Amathole and OR Tambo	2017	Taku et al. 2017
Porcine circovirus type 2	Chris Hani Amathole and OR Tambo	2017	Afolabi et al. 2017
Porcine circovirus type 2	Chris Hani Amathole and OR Tambo	2019	Afolabi et al. 2019
<i>Salmonella</i>	Amathole	2016	Iwu et al. 2016a
<i>Salmonella</i>	—	2017	Mathole et al. 2017
<i>Salmonella</i> *	OR Tambo	2019	Mthembu et al. 2019
Swine Fever	Eastern Cape**	2013	Penrith 2013
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2008	Krecek et al. 2008
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2012	Krecek et al. 2012
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2013	Krecek et al. 2013a
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2013	Krecek et al. 2013b
<i>Taenia solium</i>	Eastern Cape**	2016	Syakalime et al. 2016
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2019	Sithole et al. 2019b
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2020	Sithole et al. 2020
<i>Taenia solium</i>	OR Tambo and Alfred Nzo	2019	Sithole et al. 2019a

* – Disease found in both pigs and poultry

** – The article referred to the whole province

Table II: Poultry diseases identified in the Eastern Cape Province between 2000 and 2020 from all databases from Web of Science


Disease	Species	District	Year	Reference
Avian influenza	Ostriches	Sarah Baartman	2005	Manvell et al. 2005
Avian Influenza (H5N2)	Ostriches	Eastern Cape	2009	Abolnik et al. 2009
Avian Influenza (H5N8)	Wild birds and poultry	South Africa including Eastern Cape	2019	Abolnik 2019
<i>Salmonella</i>	Swine and chickens	OR Tambo	2019	Mthembu et al. 2019

* – Disease found in both pigs and poultry



Table III: List of pig and poultry diseases found in the Eastern Cape Province in the national database (DALRRD) from 1999 to 2019

Disease	Species	Number of reported outbreaks*
<i>Mycoplasma gallisepticum</i>	Avian	10
Newcastle disease	Avian	103
Gumboro	Avian	2
Fowl cholera	Avian	2
Avian infectious bronchitis	Avian	2
Fowl pox	Avian	18
Coccidiosis	Avian	37
<i>Salmonella enteritidis</i>	Avian	1
Avian infectious laryngotracheitis	Avian	1
Avian leukosis	Avian	9
Classical swine fever	Swine	99
Swine erysipelas	Swine	2
Cysticercosis	Swine	4
Coccidiosis	Swine	1

* – Source of data is given in Appendix 2

**Figure 2:** Frequency of poultry viral diseases reported in the Eastern Cape Province from 1999 to 2019 in the national database (DALRRD): ND – Newcastle disease: 103 outbreaks, FP – fowl pox: 18 outbreaks, AL – avian leukosis: 9 outbreaks, Gumboro and AIB – avian infectious bronchitis: 2 outbreaks, AIL – avian infectious laryngotracheitis: 1 outbreak

For pig diseases, classical swine fever (CSF) had the most reported outbreaks among pig diseases (99 outbreaks representing

**Figure 3:** Frequency of reported poultry diseases per category**Figure 4:** Frequency of reported pig diseases in the national database (DALRRD) from 1999 to 2019: classical swine fever (99 outbreaks); cysticercosis (4 outbreaks); swine erysipelas (2 outbreaks) and coccidiosis (1 outbreak).

93.4%), followed by cysticercosis (four outbreaks representing 3.8%), swine erysipelas (two outbreaks representing 1.9%) and coccidiosis (one outbreak representing 0.9%) (Figure 4).

A total number of nine diseases were retrieved from the OIE database (Table IV). The most reported poultry diseases from 2005 to 2020 were AIB and ND (reported seven times) (Table IV)

Table IV: List of pig and poultry diseases found in the OIE database (WAHID interface) from 2005 to 2020 (OIE 2020a)

Disease	Species	Number of reported outbreaks*
Fowl pox	Avian	1
Avian infectious bronchitis	Avian	7
Newcastle disease	Avian	7
LPAI (poultry)	Avian	5
HPAI	Avian	6
Gumboro	Avian	2
Mycoplasmosis	Avian	1
Fowl cholera	Avian	2
Classical swine fever	Swine	3
African swine fever	Swine	1

* – The reported outbreaks are given in detail in Appendix 3

followed by highly pathogenic avian influenza (HPAI) (reported six times). For pig diseases, the most reported disease was CSF (Table IV). Additional information on diseases prevalent in the province was obtained from the provincial laboratories despite the fact that these laboratories did not have much information on pigs and poultry diseases over the past twenty years (Appendices 4, 5 and 6).

Discussion

Data limitations

Despite the economic importance of the pig and poultry sectors in the ECP, this study found very little published information on pig and poultry diseases in the province over the past 20 years, which made it difficult to conduct a meta-analysis, as was our first intention. Also, the available published information lacked quantitative data which could help to estimate the apparent prevalence of any reported disease in the province. The national database could provide different categories of qualitative data (the status of a particular animal disease being present or absent; the species, the year in which the disease was detected, the affected area and the number of the reported cases), whereas the WAHID interface could only provide the status of the animal disease being present or absent, the species and the period (year and month) in which the disease was detected. Hence, this paper gathered information on diseases from ECP using both the national and the WAHID databases and assessed the validity of the information by comparing the findings from both.

The lack of census data in the province prevented the calculation of disease rates and comparison of years or any predictive modelling of the diseases of economic importance like ND as was performed in Zambia (Mubamba et al. 2016). These constraints limited the work presented in this paper to a descriptive review of the data available on pig and poultry diseases in the ECP but served to highlight the major deficiency in disease reporting of pig and poultry diseases in this province that has long been present.

Reporting system and the role of provincial laboratories

In the Eastern Cape, animal disease detection in village communities depends largely on the passive surveillance of pigs and poultry due to lack of human and financial resources from veterinary services. Some surveillance occurs commercially using the private laboratories outside the province, but this targets primarily the commercial sector. It is therefore likely that non-controlled diseases are not reported, especially when there is poor communication between the private sector (private veterinarians and private laboratories) and the provincial veterinary services. Active surveillance is compulsory only for export purposes (commercial farms) and is mainly for avian influenza and ND (ostriches), CSF and African swine fever (ASF) for pigs. The surveillance in the communal area (rural sector) depends on the availability of the budget and it is not done on a regular basis. The province is only equipped with three state veterinary laboratories which assist veterinary services in animal disease diagnostics and advise on the control and prevention. Unfortunately, these laboratories did not have a proper database which could be used extensively in this study. Only one laboratory could provide an electronic record of a few cases seen from 2012. It is important to mention that these provincial laboratories rely on the national laboratories for advanced diagnostic technologies, which sometimes cause a delay in finalising results and a delay in databases being updated. Private veterinarians can also send diagnostic samples directly to the national laboratory and receive results back without informing the local state veterinarians whose responsibility is to compile a comprehensive monthly report on controlled and notifiable diseases for their area. For controlled diseases, however, the accredited diagnostic veterinary laboratory is obligated by a directive to inform the state veterinarian and DALRRD directly about the results at the same time the submitter receives them.

By reviewing the references from all databases of Web of Science, it was found that the number of references reporting on diseases on the communal farms was higher (42.3%) than the number of references reporting on diseases on the commercial farms (38.5%). The references reporting on diseases on both communal and commercial farms during the same study were 11.5%, whereas three references representing 7.7% were reporting on a disease found in an abattoir. However, from the national database, it was impossible to establish whether the reported diseases were coming from the commercial or the communal farms.

By analysing the national database, the review found that ND, coccidiosis and FP were the most reported avian diseases, whereas AIB, ND and HPAI were the most reported diseases from the OIE database. For pig diseases, CSF was the most reported disease in both databases. It is suspected that these diseases were the most reported due to their outbreaks across the province in the previous years rather than active surveillance. The 2020 ASF outbreak was not found in the national database but was found on the WAHID interface database, probably because there was no update of the national database during this review, which covers the period 1993 to 2019. The review highlights the fact that the national database is less accurate in recording non-

controlled disease incidence, like Gumboro and AIB, which are known to be endemic in the province (Simbizi 2021), because it is not mandatory to report these diseases.

The limited published data, particularly on non-controlled diseases in the ECP, emphasises the need to encourage researchers to investigate animal diseases in the province.

Zoonotic diseases found in the review

A number of zoonotic diseases have been reported in the ECP. For avian influenza, a few studies identified the circulating strains (HPAI H5N2) in ostriches (Abolnik et al. 2009) and in chickens and wild birds (HPAI H5N8) (OIE 2020a). The significance of this finding in terms of human health in the province is unknown.

Despite the high number of reported cases of ND found in this study, there was no recent study investigating this disease and the circulating strains in the local poultry population. Such a study would help to understand the epidemiology of this disease for better prevention and control.

A few studies on cysticercosis (*Taenia solium*) in animals were done in the province, but they seemed to be limited to two districts (Alfred Nzo & OR Tambo) (Krecek et al. 2008; Krecek et al. 2012). This is surprising considering in 2004, an estimated 34 662 neurocysticercosis-associated cases of epilepsy were found in the ECP. The overall monetary burden (in million of US\$) was estimated to vary from US\$ 18.6 to US\$ 34.2 depending on the method used to estimate productivity losses (Carabin et al. 2006). Currently, this cost is likely to have increased given the fact that this study was done sixteen years ago. Another study on neurocysticercosis in the ECP had found that the Xhosa-speaking people of ECP had the highest prevalence of cysticercosis in South Africa probably due to the common practice of free-range pig farming and the lack of sanitation in these areas (Mafojane et al. 2003) as well as illegal slaughtering and selling of pig meats without prior meat inspection. The latter finding has been confirmed in a recent survey on trading practices of rural pig farmers in the province (Simbizi et al. 2021).

The poor sanitation in the province and the use of swine waste as manure to improve the farm yields have been mentioned as risk factors for emerging pathogens like Hepatitis E (Adelabu et al. 2017) and Norovirus (Taku et al. 2017) found in this review. Such practices will also contribute to the propagation of diseases such as *Salmonella*, *Escherichia coli*, *Campylobacter* and *Enterococcus* infections found in this review and contribute to the risk of food poisoning in rural communities of ECP. These diseases become more significant when one considers that the rate of HIV/AIDS infections in the province is among the highest in the country (Abong'o & Momba 2008).

An interesting finding was the lack of reports on diseases that one would expect to be present. Diseases like leptospirosis would have been expected to be found given the large rural pig population in the province (STATS 2016) and the fact that some serovars are maintained in pigs (Ellis 2015). With the Eastern Cape being, economically, one of the poorest provinces of South Africa, the public health impact of these neglected diseases requires further investigation.

Conclusion

This paper reviews the current knowledge on pig and poultry diseases in the rural ECP with emphasis on data from 2000 to 2020. The study found that the retrieved literature was very scarce, and little has been published on pig and poultry diseases in the ECP. Hence decision makers do not currently have reliable prior knowledge upon which to direct animal health interventions or implement public health programmes aimed at reducing the incidence of zoonotic diseases. Important neglected diseases appear not to have been studied. An improved animal health information system and further targeted research based on this study are required to fill this gap in knowledge.

Poor communication between important disease reporting stakeholders in the province was reflected in the review through disparities in data sources and it is recommended that this be improved. Improved communication between the national department (DALRRD) and the National Institute of Communicable Diseases will increase awareness about the zoonotic diseases found in this review and help to minimise their impact on the livelihoods of the rural communities. It is recommended, therefore, that a disease reporting system in the province involving all the stakeholders be considered to provide current relevant information on pig and poultry diseases. This will provide a foundation for sound decision making around disease control and international trade in live animals and animal products.

Author contributions

V Simbizi conducted the literature review, analysed and interpreted the data under the supervision of B Gummow. V Simbizi wrote the article under the supervision of B Gummow. All authors assisted in the editing and critiquing of the manuscript.

Acknowledgements

Grahamstown Provincial Veterinary Laboratory for allowing access to the laboratory records; Middleburg Provincial Veterinary Laboratory for allowing access to the laboratory records.

Conflict of interest

The authors declare that no conflict of interest.

Funding source

No funding was required.

Data availability statement

A list of figures and tables that have associated raw data are included as appendices.

Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors.

Ethical approval

This article followed all ethical standards for carrying out research without direct contact with human or animal subjects.

ORCID

V Simbizi  <https://orcid.org/0000-0003-0337-4721>

R Moerane  <https://orcid.org/0000-0002-7711-6189>
 G Ramsay  <https://orcid.org/0000-0003-0651-9308>
 C Mubamba  <https://orcid.org/0000-0003-3669-1299>
 C Abolnik  <https://orcid.org/0000-0003-3044-465X>
 B Gummow  <https://orcid.org/0000-0001-6364-1583>

References

- Abolnik, C., 2019, Outbreaks of Clade 2.3.4.4 H5N8 highly pathogenic avian influenza in 2018 in the northern regions of South Africa were unrelated to those of 2017, *Transbound Emerg Dis* 67(3), 1371–1381. <https://doi.org/10.1111/tbed.13448>.
- Abolnik, C., Londt, B.Z., Manvell, R.J., et al., 2009, Characterisation of a highly pathogenic influenza A virus of subtype H5N2 isolated from ostriches in South Africa in 2004, *Influenza Other Respir Viruses* 3, 63–68. <https://doi.org/10.1111/j.1750-2659.2009.00074.x>.
- Abong'o, B.O., Momba, M.N.B., 2008, Prevalence and potential link between *E. coli* O157 : H7 isolated from drinking water, meat and vegetables and stools of diarrhoeic confirmed and non-confirmed HIV/AIDS patients in the Amathole District – South Africa, *J Appl Microbiol* 105, 424–431. <https://doi.org/10.1111/j.1365-2672.2008.03756.x>.
- Adelabu, O.A., Iweriebor, B.C., Nwodo, U.U., et al., 2017, Incidence and molecular characterization of hepatitis E virus from swine in Eastern Cape, South Africa, *Adv Virol* 2017, 1073253. <https://doi.org/10.1155/2017/1073253>.
- Afolabi, K.O., Iweriebor, B.C., Obi, L.C., Okoh, A.I., 2017, Molecular detection of Porcine circovirus type 2 in swine herds of Eastern Cape Province South Africa, *BMC Microbiol* 17, 212. <https://doi.org/10.1186/s12866-017-1121-4>.
- Afolabi, K.O., Iweriebor, B.C., Obi, C.L., Okoh, A.I., 2019, Genetic characterisation and diversity of porcine circovirus type 2 in non-vaccinated South African swine herds, *Transbound Emerg Dis* 66, 412–421. <https://doi.org/10.1111/tbed.13036>.
- Akol, G.W., Lubisi, B.A., 2010, Classical swine fever control in South Africa 2008–09: results of the disease surveillance in the Eastern Cape Province, Pretoria, South Africa, *Southern African Society for Veterinary Epidemiology and Preventive Medicine*.
- Carabin, H., Krecek, R.C., Cowan, L.D., et al. 2006. Estimation of the cost of *Taenia solium* cysticercosis in Eastern Cape Province, South Africa, *Trop Med Int Health* 11, 906–916. <https://doi.org/10.1111/j.1365-3156.2006.01627.x>.
- DAFF 2016. Animal Disease Reporting Manual.
- DAFF 2018. A profile of the South African Pork Market Value Chain. Available from: <https://www.nda.agric.za/doiDev/sideMenu/Marketing/Annual%20Publications/Commodity%20Profiles/Pork%20Market%20Value%20Chain%20Profile%202018.pdf>. Accessed 4 Mar 2020.
- DAFF 2020. Query on Animal Diseases in the RSA. Available from: <http://webapps.daff.gov.za/VetWeb/diseaseDatabase.do>. Accessed 1 Feb 2020.
- ECSECC 2018. Economic review of the Eastern Cape Gross Domestic Product (GDP) Quarter 2-2018. Available from: https://www.ecsecc.org/documentrepository/informationcentre/ecsecc-gdp-report-oct-2018final_08658.pdf. Accessed 22 Aug 2020.
- Ellis, W.A., 2015, Animal Leptospirosis, in B. Adler, (ed.) *Leptospira and Leptospirosis*. Berlin: Springer-Verlag Berlin. https://doi.org/10.1007/978-3-662-45059-8_6.
- FAO 2020. Transboundary animal diseases. FAO. Available from: <http://www.fao.org/emergencies/emergency-types/transboundary-animal-diseases/en/>. Accessed 19 Jan 2020.
- Igwaran, A., Okoh, A., 2020, Campylobacteriosis Agents in meat carcasses collected from two district municipalities in the Eastern Cape Province, South Africa, *Foods* 9(2), 203. <https://doi.org/10.3390/foods9020203>.
- Iweriebor, B.C., Obi, L.C., Okoh, A.I., 2015, Virulence and antimicrobial resistance factors of *Enterococcus* spp. isolated from fecal samples from piggery farms in Eastern Cape, South Africa, *BMC Microbiol* 15, (4 July 2015). <https://doi.org/10.1186/s12866-015-0468-7>.
- Iwu, C.J., Iweriebor, B.C., Obi, L.C., et al., 2016a, Multidrug-resistant *Salmonella* isolates from swine in the Eastern Cape Province, South Africa, *J Food Prot* 79, 1234–1239. <https://doi.org/10.4315/0362-028X.JFP-15-224>.
- Iwu, C.J., Iweriebor, B.C., Obi, L.C., Okoh, A.I., 2016b, Occurrence of non-O157 Shiga toxin-producing *Escherichia coli* in two commercial swine farms in the Eastern Cape Province, South Africa, *Comp Immunol Microbiol Infect Dis* 44, 48–53. <https://doi.org/10.1016/j.cimid.2015.12.004>.
- Iwu, C.J., Jaja, I.F., Iweriebor, B.C., et al., 2017, Antibiotic resistance profiles of *Escherichia coli* O26, O145, and O157:H7 isolated from swine in the Eastern Cape Province, South Africa, *Asian Pac J Trop Dis* 7, 553–559. <https://doi.org/10.12980/apjtd.7.2017D7-9>.
- Krecek, R.C., Michael, L.M., Schantz, P.M., et al., 2008, Prevalence of *Taenia solium* cysticercosis in swine from a community-based study in 21 villages of the Eastern Cape Province, South Africa, *Vet Parasitol* 154, 38–47. <https://doi.org/10.1016/j.vetpar.2008.03.005>.
- Krecek, R.C., Mohammed, H., Michael, L.M., et al., 2012, Risk factors of porcine cysticercosis in the Eastern Cape Province, South Africa, *PLoS One* 7, e37718. <https://doi.org/10.1371/journal.pone.0037718>.
- Krecek, R.C., Mohammed, H., Michael, L.M., et al., 2013a, Bivariable associations between owner/pig characteristics and cysticercosis1 infection in pigs from Eastern Cape Province (South Africa) (N=256)2. Figshare, *PLoS One*. Dataset. <https://doi.org/10.1371/journal.pone.0037718.t001>.
- Krecek, R.C., Mohammed, H., Michael, L.M., et al., 2013b, Final multivariable model for the association between owner/pig characteristics and cysticercosis1 infection in pigs from Eastern Cape Province (South Africa) (N=256)2. Figshare, *PLoS One*. Dataset. <https://doi.org/10.1371/journal.pone.0037718.t002>.
- Mafojane, N.A., Appleton, C.C., Krecek, R.C., et al., 2003, The current status of neurocysticercosis in Eastern and Southern Africa, *Acta Trop* 87, 25–33. [https://doi.org/10.1016/S0001-706X\(03\)00052-4](https://doi.org/10.1016/S0001-706X(03)00052-4).
- Manvell, R.J., Horner, R., Akol, G., et al., 2005, Isolation of an influenza A virus subtype H5N2 from ostriches in South Africa in 2004.
- Mathole, M.A., Muchadeyi, F.C., Mdladla, K., et al., 2017, Presence, distribution, serotypes and antimicrobial resistance profiles of *Salmonella* among pigs, chickens and goats in South Africa, *Food Cont* 72, 219–224. <https://doi.org/10.1016/j.foodcont.2016.05.006>.
- Morris, R.S., 1991, Information systems for animal health: objectives and components, *Rev Sci Tech* 10, 13–23. <https://doi.org/10.20506/rst.10.1.537>.
- Mthembu, T.P., Zishiri, O.T., El Zowalaty, M.E., 2019, Detection and molecular identification of *Salmonella* virulence genes in livestock production systems in South Africa, *Pathogens* 8, 124. <https://doi.org/10.3390/pathogens8030124>.
- Mubamba, C., Ramsay, G., Abolnik, et al., 2016, A retrospective study and predictive modelling of Newcastle Disease trends among rural poultry of eastern Zambia, *Prev Vet Med* 133, 97–107. <https://doi.org/10.1016/j.prevetmed.2016.09.017>.
- Ngobese, B., Zishiri, O.T., El Zowalaty, M.E. 2020. Molecular detection of virulence genes in *Campylobacter* species isolated from livestock production systems in South Africa, *J Integr Agric* 19, 1656–1670. [https://doi.org/10.1016/S2095-3119\(19\)62844-3](https://doi.org/10.1016/S2095-3119(19)62844-3).
- OIE 2020a. WAHIS Interface. Available from: <https://www.oie.int/en/?s=&search=WAHIS+Interface>. Accessed 1 Feb 2020.
- OIE 2020b. World Animal Health Information Database (WAHIS) Interface. OIE, Available from: https://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home. Accessed 21 Jan 2020.
- Penrith, M.L., 2013, History of 'swine fever' in Southern Africa. *J S Afr Vet Assoc* 84, 1106. <https://doi.org/10.4102/jsava.v84i1.1106>.
- SAPA 2017. Distribution of chickens in South Africa for the surveillance period July 2017 to December 2017. Available from: <http://www.sapoultry.co.za/pdf-statistics/provisional-distribution-of-chickens-in-sa.pdf>. Accessed 4 Mar 2020.
- Simbizi, V., Moerane, R., Ramsay, G., et al., 2021. A study of rural chicken farmers, diseases and remedies in the Eastern Cape Province of South Africa, *Prev Vet Med* 194, 105430. <https://doi.org/10.1016/j.prevetmed.2021.105430>.
- Sithole, M. I., Bekker, J. L., Mukaratirwa, S., 2019a, Pig husbandry and health practices of farmers in selected *Taenia solium* endemic rural villages of two districts in the Eastern Cape Province of South Africa, *Int J Vet Sci* 8, 235–242.
- Sithole, M.I., Bekker, J.L., Mukaratirwa, S., 2020, Consumer knowledge and practices to pork safety in two *Taenia solium* cysticercosis endemic districts in Eastern Cape Province of South Africa, *BMC Infect Dis*, 20, 107. <https://doi.org/10.1186/s12879-020-4839-9>.
- Sithole, M.I., Bekker, J.L., Tsotetsi-Khambule, A.M., Mukaratirwa, S., 2019b, Ineffectiveness of meat inspection in the detection of *Taenia solium* cysticerci in pigs slaughtered at two abattoirs in the Eastern Cape Province of South Africa, *Vet Parasitol Reg Stud Rep* 17, 100299. <https://doi.org/10.1016/j.vprsr.2019.100299>.
- STATS 2016. Community Survey 2016 Agricultural households. Statistics South Africa.
- STATS 2018. Gross Domestic Product 4th quarter 2018. Available from: http://www.statssa.gov.za/publications/P0441/GDP_2018_Q4_Media_presentation.pdf#page=12. Accessed 22 Aug 2020.
- STATS 2020. Mid-year population estimates 2020. Available from: <http://www.statssa.gov.za/publications/P0302/P03022020.pdf>. Accessed 23 Aug 2020.
- Syakalime, M., Foli, T.L., Mwanza, M., 2016, Risk factors and prevalence of Porcine cysticercosis in free range pigs of selected areas of South Africa, *Indian Journal of Animal Research* 50, 287–289. <https://doi.org/10.18805/ijar.8420>.
- Taku, O., Iweriebor, B.C., Nwodo, U.U., et al., 2017, Occurrence of Norovirus in pig faecal samples in the Eastern Cape, South Africa, *Asian Pac J Trop Dis* 7, 151–155. <https://doi.org/10.12980/apjtd.7.2017D6-393>.